

ON FLEXIBLE QUADRATIC ALGEBRAS

S. PUMPLÜN

ABSTRACT. Let R be a ring. A construction method for flexible quadratic algebras with scalar involution over R is presented which unifies various classical constructions in the literature, in particular those to construct composition algebras.

INTRODUCTION

Let R be a unital commutative associative ring. We present a construction method for R -algebras starting from an associative quadratic R -algebra with a scalar involution. Special cases of this construction yield flexible quadratic algebras over rings (an algebra A is *flexible* if $x(yx) = (xy)x$ for all elements $x, y \in A$). Composition algebras are certainly among the best-known classes of flexible quadratic algebras. Our method unifies the different approaches by Petersson [P2], Thakur [T] and [Pu] to construct composition algebras. In special cases it overlaps with the one described by Becker [B] which yields flexible quadratic division algebras containing subalgebras of half their rank.

1. PRELIMINARIES

Let A be an R -algebra. The term “ R -algebra” always refers to a unital nonassociative algebra A which is faithfully projective as an R -module [K, p. 52].

A is called *alternative* if its associator $[x, y, z] = (xy)z - x(yz)$ is alternating. The *nucleus* of A is defined as $N(A) = \{x \in A \mid [x, A, A] = [A, x, A] = [A, A, x] = 0\}$. The nucleus is an associative subalgebra of A (it may be zero), and $x(yz) = (xy)z$ whenever one of the elements x, y, z is in $N(A)$. An anti-automorphism $\sigma : A \rightarrow A$ of period 2 is called an *involution* on A . If 2 is an invertible element in R , we have $A = \text{Sym}(A, \sigma) \oplus \text{Skew}(A, \sigma)$ with $\text{Skew}(A, \sigma) = \{x \in A \mid \sigma(x) = -x\}$ the set of skew symmetric elements

2000 *Mathematics Subject Classification*. Primary: 17A45; Secondary: 17A20, 17A75, 11E25.

Key words and phrases. Flexible quadratic algebras, composition algebras, cross products.

and $\text{Sym}(A, \sigma) = \{x \in A \mid \sigma(x) = x\}$ the set of symmetric elements in A with respect to σ . An involution is called *scalar* if all *norms* $\sigma(x)x$ are elements of R . For every scalar involution σ , $n_A(x) = \sigma(x)x$ (resp. the *trace* $t_A(x) = \sigma(x) + x$) is a quadratic (resp. a linear) form on A . A is called *quadratic*, if there exists a quadratic form $n: A \rightarrow R$ such that $n(1_A) = 1$ and $x^2 - n(1_A, x)x + n(x)1_A = 0$ for all $x \in A$, where $n(x, y)$ denotes the induced symmetric bilinear form $n(x, y) = n(x + y) - n(x) - n(y)$. The form n is uniquely determined and called the *norm* of the quadratic algebra A [P2]. The existence of a scalar involution on an algebra A implies that A is quadratic [M1].

An R -algebra C is called a *composition algebra* if it carries a quadratic form $n: C \rightarrow R$ whose induced symmetric bilinear form $n(x, y)$ is *nondegenerate* (i.e., determines an R -module isomorphism $C \xrightarrow{\sim} C^\vee = \text{Hom}_R(C, R)$) and which satisfies $n(xy) = n(x)n(y)$ for all $x, y \in C$ (it *permits composition*). Composition algebras are quadratic alternative; any nondegenerate quadratic form n on the composition algebra which permits composition is its norm as a quadratic algebra and thus is unique [P2]. It is called the *norm* of C and is also denoted by n_C . A quadratic alternative algebra is a composition algebra if and only if its norm is nondegenerate [M1, 4.6]. Composition algebras only exist in ranks 1, 2, 4 or 8. Those of rank 2 are exactly the quadratic étale R -algebras, those of rank 4 exactly the well-known quaternion algebras. The ones of rank 8 are called *octonion algebras*. A composition algebra C has a *canonical involution* $\bar{}$ given by $\bar{x} = t_C(x)1_C - x$, where $t_C: C \rightarrow R$, $t_C(x) = n(1_C, x)$, is the *trace* of C . This involution is scalar.

2. FLEXIBLE QUADRATIC ALGEBRAS

Let N be a finitely generated projective R -module of constant rank. An alternating R -bilinear map $\times: N \times N \rightarrow N$ is called a *cross product* on N . Let D be an associative R -algebra with a scalar involution $\sigma = \bar{}$ (e.g. a quadratic étale or a quaternion algebra over R with canonical involution). Let F be a locally free right D -module of constant finite rank s together with a sesquilinear form $h: F \times F \rightarrow D$ (i.e., h is a biadditive map such that $h(ua, vb) = \bar{a}h(u, v)b$ for all $a, b \in D$, $u, v \in F$). Let $\times$ be a cross product on F , where now F is viewed as an R -module, i.e. together with $\times: F \times F \rightarrow F$, F is an anticommutative R -algebra. The R -module $A = D \oplus F$ becomes a unital R -algebra denoted by $A = (D, F, h, \times)$ via the multiplication

$$(\alpha, u)(\beta, v) = (\alpha\beta - h(v, u), v\alpha + u\bar{\beta} + v \times u),$$

for all $\alpha, \beta \in D$, $u, v \in F$. The algebra D is a subalgebra of $(D, F, h, \times)$.

Define

$$\begin{aligned}\sigma_A : A &\rightarrow A, & (\alpha, u) &\rightarrow (\bar{\alpha}, -u), \\ n_A : A &\rightarrow D, & n_A((\alpha, u)) &= \sigma_A(\alpha, u)(\alpha, u), \\ t_A : A &\rightarrow R, & t_A((\alpha, u)) &= \sigma_A(\alpha, u) + (\alpha, u) = (t_D(\alpha), 0).\end{aligned}$$

Obviously, σ_A is an R -module isomorphism of order 2. We have

$$\begin{aligned}n_A((\alpha, u)) &= n_D(\alpha) + h(u, u), \\ n_A((\alpha, u)) &= (\alpha, u)\sigma_A(\alpha, u), \\ n_A((\sigma(\alpha, u))) &= n_A((\alpha, u)) \text{ and } t_A(\sigma(\alpha, u)) = t_A((\alpha, u)).\end{aligned}$$

The map n_A is quadratic with associated R -bilinear map $n_A : A \times A \rightarrow D$ given by

$$\begin{aligned}n_A((\alpha, u), (\beta, v)) &= n_D(\alpha, \beta) + (h(u, v) + h(v, u)) = \\ &= (\alpha, u)\sigma_A(\beta, v) + (\beta, v)\sigma_A(\alpha, u).\end{aligned}$$

Obviously, $F = \ker(t_A) = \ker(t_D) \oplus F$ and $u \times v = uv - \frac{1}{2}n_A(u, v)$.

Lemma 1. (i) σ_A is a scalar involution if and only if h is a hermitian form (i.e., $h(u, v) = \overline{h(v, u)}$ for all $u, v \in F$).
(ii) If h is a hermitian form, then $t : A \times A \rightarrow R$, $t(x, y) = t_A(xy)$ is a symmetric R -bilinear form.

Lemma 1 is a weak generalization of [B, Lemma 1].

Corollary 1. If $h : F \times F \rightarrow D$ is a (perhaps degenerate) hermitian form, then $A = (D, F, h, \times)$ is a quadratic R -algebra with scalar involution $\sigma_A : A \rightarrow A$, $\sigma_A((\alpha, u)) = (\bar{\alpha}, -u)$ and norm $n_A : A \rightarrow R$, $x \rightarrow x\sigma_A(x)$, where $n_A((\alpha, u)) = n_D(\alpha) + h(u, u)$. Moreover, if n_A is isotropic (e.g., if h is isotropic) then A has zero divisors.

Note that even if D is an associative composition division algebra over a field, the fact that two quadratic algebras $A = (D, F, h, \times)$ and $A' = (D, F', h', \times')$ have isometric norms $n_A \cong n_{A'}$ only implies that the non-degenerate hermitian forms (F, h) and (F', h') are isometric and yields no information on the cross products $\times$ and $\times'$.

Quadratic R -algebras with a scalar involution were constructed using finite dimensional vector spaces which carry a bilinear form B by Osborn [O]. Finite-dimensional quadratic division algebras over a field of characteristic not 2 were shown to be even-dimensional by Petersson [P1]. Already earlier, Kuz'min [Ku] proved that any quadratic division algebra, over a field of

characteristic not 2 in which any two elements generate a subalgebra of dimension at most four, must have finite dimension and that, indeed, only the dimensions 1, 2, 4 or 8 are possible. Independently, Kunze and Scheinberg [KS, Theorem 2.4] enumerated all unital alternative algebras over fields of characteristic not 2 - including the infinite-dimensional ones - which have a scalar involution.

Remark 1. For $A = (D, F, h, \times)$, $2 \in R^\times$, we obtain $\text{Sym}(A, \sigma_A) = R$ and $\text{Skew}(A, \sigma_A) = \text{Skew}(D, \sigma_D) \oplus F$.

Lemma 2. *If $h : F \times F \rightarrow D$ is a hermitian form, then $(D, F, h, \times)$ is*
(i) flexible if and only if $h(u \times v, u) + \overline{h(u \times v, u)} = n_A(u \times v, u) = 0$ and $(u \times v) \times u = u \times (v \times u)$ for all $u, v \in F$;
(ii) alternative if and only if $h(u, u \times v) = 0$ and $u \times (u \times v) = -h(u, u)v + h(v, u)u$ for all $u, v \in F$, if and only if $h(u \times v, v) = 0$ and $(u \times v) \times v = h(v, v)u - h(u, v)v$ for all $u, v \in F$.

If $(D, F, h, \times)$ is flexible with a scalar involution, then it is a noncommutative Jordan algebra [M1, (3.3)]. Due to the nature of the multiplication in $(D, F, h, \times)$ it suffices to work with elements of F to check flexibility. The identities in (ii) were already obtained in [T, (2.1)] in a more restricted setting (there, F has rank 3, D is quadratic étale, and h is a nondegenerate hermitian form). If $\times$ is the zero-map, then $(D, F, h, 0)$ is trivially flexible.

Remark 2. If $D = R$, then $\sigma = id$ and $h = B$ is a symmetric R -bilinear form on F . $A = (R, F, B, \times)$ is the unital quadratic R -algebra described in [B, p. 26] (see also [O, Theorem 1]). It is well-known that any quadratic algebra A over a field of characteristic not 2 with a scalar involution can be of the type $A = (R, F, B, \times)$ for a suitable F , a symmetric bilinear form B and a cross product $\times$.

(i) By Lemma 2 (i), A is flexible if and only if $B(u, v \times u) = B(u, u \times v)$ and $(u \times v) \times u = u \times (v \times u)$ for all $u, v \in F$, which is equivalent to

$$B(u, v \times u) = 0 \quad \text{and} \quad (u \times v) \times u = u \times (v \times u)$$

for all $u, v \in F$.

(ii) Let $2 \in R^\times$. If we apply [B, Lemma 2, Satz 1] to the residue class algebras, we see that A is flexible if and only if t_A is associative, if and only if B is associative; and that A is flexible and t_A and n_A are weakly nondegenerate if and only if B is associative and weakly nondegenerate. (q is *weakly nondegenerate* if $\text{Rad}(q) = \{a \in A \mid q(a, A) = 0\} = 0$.)

(iii) For a flexible quadratic R -algebra $A = (R, F, B, \times)$ of rank greater or equal to 3 with a weakly nondegenerate associative bilinear form B , the residue class algebra $A(P)$ is central-simple for each $P \in \text{Spec}(R)$. This follows from [B, Satz 1, Satz 2] applied to the residue class algebras. If A is a quadratic algebra of rank 3 over a domain R and B weakly nondegenerate and associative, then A is commutative. This follows from [B, p. 33].

Theorem 1. *Let $2 \in R^\times$.*

(i) Let F be a finitely generated projective R -module of constant rank s . Let $B : F \times F \rightarrow R$ be a nondegenerate symmetric bilinear form on F and $\times : F \times F \rightarrow F$ a cross product, such that

$$(1) \quad B(u \times v, u \times v) = B(u, u)B(v, v) - B(u, v)^2$$

$$(2) \quad B(u, u \times v) = 0$$

for all $u, v \in F$. Then $C = (R, F, B, \times)$ is a composition algebra over R . In particular $s = 1$, $s = 3$, or $s = 7$, and $\det(F, N) = \bigwedge^s(F, N) = R$.

(ii) Let C be a composition algebra over R with norm n of rank $s + 1 \geq 1$. Then $C = R \oplus F$ with $F = \text{Skew}(C, \bar{})$. The R -module F satisfies $\det(F) = \bigwedge^s(F) = R$ with $s = 1, 3$ or 7 . Define $\times : F \times F \rightarrow F$ via $u \times v = \frac{1}{2}[u, v] = \frac{1}{2}(uv - vu)$ and $B : F \times F \rightarrow R$ via $B(u, v) = -\frac{1}{2}(uv + vu) = \frac{1}{2}(u\bar{v} + v\bar{u}) = \frac{1}{2}n(u, v)$. Then $\times$ is a cross product and B is a nondegenerate symmetric bilinear form on F satisfying equations (1) and (2) in (i) for all $u, v \in F$.

This is well-known if R is a field. Lemma 2 (ii) implies the first part of Theorem 1. Otherwise, the proof of [Mo, Theorem 10] can be used verbatim. Note that the intrinsic value of this theorem is impaired by the fact that producing a cross product on a finitely generated projective R -module F of rank 7 satisfying (1) and (2) is probably not any easier than producing an octonion algebra structure on $R \oplus F$.

If, in (ii), C is quadratic étale, then there exists a line bundle L over R and a nondegenerate quadratic form $N_L : L \times L \rightarrow R$ such that $C \cong \text{Cay}(R, L, N_L)$. Since every bilinear map $L \times L \rightarrow L$ is symmetric, this forces the cross product on L to be zero.

Example 1. Let A be a quadratic R -algebra of rank 2. Then A is commutative associative and the conjugation is a scalar involution (see [K, I.(1.3.6)] or check it locally). We get $A = (R, L, B, \times)$ for a suitable $L \in \text{Pic } R$,

$B : L \times L \rightarrow R$ a bilinear form. A is commutative, therefore $\times$ is the zero-map, and $A = \text{Cay}(R, L, N)$ with $N : L \rightarrow R$ the - perhaps degenerate - quadratic form associated to B .

Lemma 3. *Let $2 \in R^\times$. Let A be a flexible quadratic R -algebra with a scalar involution σ which contains a composition subalgebra D of rank 2 and no other associative composition subalgebra of larger rank, or a composition subalgebra D of rank 4. Let $F = D^\perp$. Define $-h$ to be the orthogonal projection of the multiplication vu for $u, v \in F$ onto D and $\times$ to be the orthogonal projection of the multiplication vu for $u, v \in F$ onto F . Then:*

- (i) $\times : F \times F \rightarrow F$ is a cross product.
- (ii) $n_A(u, v) = t_D(h(u, v))$ and $h(u, v) = \overline{h(v, u)}$.

Proof. Let A be a quadratic R -algebra with a scalar involution σ , which contains a composition subalgebra D of rank 2 or 4 as subalgebra of maximal rank with nondegenerate norm. Since the norm $n_A(x) = x\sigma(x)$ is nondegenerate when restricted to D , we can decompose A as $A = D \perp D^\perp$ [K, (3.6.2), p. 17]. Since $-h$ and $\times$ are the orthogonal projections of the multiplication vu for $u, v \in F$ onto D and onto F , we have $vu = -h(u, v) + u \times v$ for $u, v \in F$.

- (i) Since $F \subset \text{Skew}(A, \sigma)$, it follows that $u^2 \in R$ for all $u \in F$ and thus $u \times u = u^2 + h(u, u) \in D \cap D^\perp = \{0\}$. Hence $\times$ is anticommutative and a cross product on F .
- (ii) We have $n_A(u) = -u^2 = h(u, u)$ for all $u \in F$. Since $u \times v = -v \times u$, it follows that $h(u, v) + \overline{h(u, v)} = -vu - uv = v\sigma(u) + u\sigma(v) = n_A(u, v)$, i.e. $n_A(u, v) = t_D(h(u, v))$ and $h(u, v) = \overline{h(v, u)}$. $\square$

Example 2. (a) Let $2 \in R^\times$. Let $A = (D, F, h, \times)$ be an algebra with a scalar involution, D quadratic étale or a quaternion algebra. The algebra $\hat{A} = (R, F, B, \times)$, $B(u, v) = \frac{1}{2}(h(u, v) + \overline{h(u, v)}) = \frac{1}{2}n_A(u, v)$, is flexible if and only if A is flexible. If A is alternative, the identity $((u \times v) \times v) \times v = n_A(v)u \times v$ holds in $\hat{A}$ for all $u, v \in F$ (Lemma 2). In particular, if $A = \text{Cay}(T, F, h, \times_\alpha)$ is an octonion algebra over a field R of characteristic not 2, then $\hat{A} = (R, F, B, \times_\alpha)$ is the colour algebra defined in [E].

(b) If the cross product is trivial, $A = (D, F, h, 0)$ with S a quadratic étale R -algebra or a quaternion algebra over R , is a unital noncommutative Jordan algebra. In particular, $(R, F, -B, 0) = \text{JSpin}(F, B)$ is the well-known algebra of the Spin Factors [M2, p. 178].

(c) Let k be a field of characteristic not 2. Given a quadratic k -algebra A

with scalar involution $*$, the map $(a, b)^* = (a^*, -b)$ defines a scalar involution on the quadratic algebra $C(A, \cdot_1, \cdot_2, \cdot_3) = A \oplus A$ with multiplication $(a, b)(c, d) = (ac + d^* \cdot_1 b, d \cdot_2 a + b \cdot_2 c^* + b \cdot_3 d)$, where the maps $\cdot_i$ are certain bilinear products on A [B]. Using this construction all flexible quadratic division algebras of dimension $2m$ over k which contain a unital subalgebra of dimension m can be obtained [B, p. 41]. The construction generalizes to quadratic algebras with scalar involutions over arbitrary rings where 2 is invertible. However, it will not produce all such algebras. If a flexible quadratic algebra over a field contains a composition subalgebra of dimension 2 or 4, the method coincides with ours whenever the algebra has twice its dimension, i.e. dimension 4, respectively 8. In that case $d^* \cdot_1 b$ is given by the hermitian form $h(d, b)$, $\cdot_2$ is the usual algebra multiplication of A , and $\cdot_3$ is our cross product $\times$.

Let S be a quadratic étale R -algebra. For the remainder of this section, we restrict our attention to the special case that F is a finitely generated projective right S -module of constant rank carrying a nondegenerate hermitian form $h: F \times F \rightarrow S$. For each invertible $\mu \in S$, the form $\mu h: F \times F \rightarrow S$ is a nondegenerate ϵ -hermitian form, with $\epsilon = \bar{\mu}/\mu$. In particular, $\epsilon \neq 1$ if and only if $\mu \notin R$, and $\epsilon = -1$ if and only if $\mu \in \text{Skew}(S, \bar{})$.

Recall that conversely, the study of ϵ -hermitian forms over S can be reduced to the study of hermitian ones by scaling in several important cases: Let $\mu \in S^\times$ such that $\frac{\bar{\mu}}{\mu} = \epsilon$. Then μh is a hermitian form, for any ϵ -hermitian form h . Such a μ exists if $H^1(\mathbb{Z}/2\mathbb{Z}, S^\times) = 0$ (Hilbert's Theorem 90), e.g. if $\text{Pic } R = 0$ [K, p. 300].

Proposition 1. *Let $A = (S, F, \mu h, \times)$ for some nondegenerate hermitian form $h: F \times F \rightarrow S$, and an invertible scalar $\mu \in S \setminus R$. If S is a domain (or if F is torsion-free and $\bar{\mu} - \mu$ not a zero divisor in S), then $N(A) = S$.*

Proof. A straightforward calculation shows that $S \subset N(A)$.

Now let $(e, e') \in N(A)$ with $e \in S$ and $e' \in F$. We have to show that this implies $e' = 0$. The equation

$$(e, e')((u, w)(u', w')) = ((e, e')(u, w))(u', w')$$

implies that

$$e' \bar{\mu} h(w, w') = w' \mu h(w, e')$$

for all $w, w' \in F$ and thus (put $w' = e'$)

$$e'(\bar{\mu} - \mu)h(w, e') = 0$$

for all $w \in F$. If $h(w, e') = 0$ for all $w \in F$ then $e' = 0$ since h is nondegenerate and thus $(e, e') = (e, 0) \in S$. Otherwise, there is one $w \in F$ such that $h(w, e') \neq 0$. Then if S has no zero divisors, i.e., is a domain, F is torsion free and we have $e' = 0$ since $\bar{\mu} - \mu \neq 0$ here. The same holds if F is torsion free and $\bar{\mu} - \mu$ not a zero divisor in S . $\square$

3. CONSTRUCTIONS OF COMPOSITION ALGEBRAS

Petersson [P2] introduced the following *generalized Cayley-Dickson doubling process* for composition algebras over rings: Let D be a composition algebra of rank ≤ 4 over R with canonical involution $\bar{\cdot}$. Let P be a locally free right D -module of rank one and norm one (cf. [P2, 2.3] for the definition; if D is a quaternion algebra, of norm one means that the reduced norm functor defined in [KOS] associates to P the free line bundle R). Thus there exists a nondegenerate quadratic form $N: P \rightarrow R$ satisfying $N(w \cdot u) = N(w)N_D(u)$ for all elements $w \in P$, $u \in D$, where $\cdot$ denotes the right D -module structure of P (for D a quaternion algebra, see also [K, p. 161 ff]). N is uniquely determined up to an invertible factor in $R^\times$ and determines a unique R -bilinear map $P \times P \rightarrow D$, written multiplicatively and satisfying $(w \cdot u)(w \cdot v) = N(w)\bar{v}u$ for all $u, v \in D$, $w \in P$. The R -module $\text{Cay}(D, P, N) = D \oplus P$ becomes a composition algebra under the multiplication

$$(u, w)(u', w') = (uu' + ww', w' \cdot u + w \cdot \bar{u}')$$

for $u, u' \in D$, $w, w' \in P$, with $\cdot$ denoting the right D -module structure of P . Its norm is given by

$$n_{\text{Cay}(D, P, N)}((u, w)) = n_D(u) - N(w).$$

If F a locally free right D -module of rank 1 and $h: F \times F \rightarrow D$ a nondegenerate hermitian form, then

$$(D, F, h, 0) = \text{Cay}(D, F, -N)$$

where $N(u) = h(u, u)$ for all $u \in F$.

Let S be a quadratic étale R -algebra with canonical involution $\bar{\cdot}$. Let (P, h) be a ternary nondegenerate $\bar{\cdot}$ -hermitian space such that $\bigwedge^3(P, h) \cong \langle 1 \rangle$. Choose an isomorphism $\alpha: \bigwedge^3(P, h) \rightarrow \langle 1 \rangle$. Define a cross product $\times_\alpha: P \times P \rightarrow P$ as in [T, p. 5122]. Note that $\times_\alpha$ can be derived solely from T, P, h and α . The R -module $\text{Cay}(S, P, h, \alpha) = S \oplus P$ becomes an octonion

algebra under the multiplication

$$(a, u)(b, v) = (ab - h(v, u), va + u\bar{b} + u \times_\alpha v)$$

for $u, v \in P$ and $a, b \in S$, with norm

$$n((a, u)) = n_S(a) + h(u, u).$$

If the ternary hermitian space (P, h) is orthogonally decomposable this construction is independent of the choice of the isomorphism α . Any octonion algebra which contains a quadratic étale subalgebra can be constructed like this (Petersson-Racine [PR, 3.8] or Thakur [T]). We obtain

$$(S, P, h, \times_\alpha) = \text{Cay}(S, P, h, \alpha).$$

Let $2 \in R^\times$. There exists a general construction for quaternion algebras over R [Pu]: Let F be a finitely generated projective R -module of constant rank 3 such that $\det F = \bigwedge^3 F \cong R$, and $N: F \rightarrow R$ a nondegenerate quadratic form, i.e., a selfdual isomorphism $\varphi_N: F \xrightarrow{\sim} \check{F}$, $x \rightarrow N(x, _)$. (Any selfdual isomorphism φ from F to its dual bundle $\check{F}$ canonically induces a nondegenerate quadratic form on F via $N_\varphi(x) := \frac{1}{2}\langle x, \varphi(x) \rangle$ where $\langle _, _ \rangle: F \times \check{F} \rightarrow R$ is the canonical pairing, cf. [P2, 3.2].) Assume that N has trivial determinant. Choose an isomorphism $\alpha: \det F \xrightarrow{\sim} R$, then α is unique up to an invertible factor in R and induces a bilinear map

$$F \times F \rightarrow \check{F}$$

$$(u, v) \rightarrow u \times_\alpha v := \alpha(u \wedge v \wedge _)$$

as well as an isomorphism $\beta: \det \check{F} \xrightarrow{\sim} R$ given by the condition

$$\alpha(u_1 \wedge u_2 \wedge u) \beta(\check{u}_1 \wedge \check{u}_2 \wedge \check{u}_3) = \det(\langle u_i, \check{u}_j \rangle)$$

for u_i in F , $\check{u}_j$ in $\check{F}$, $1 \leq i, j \leq 3$, where again $\langle _, _ \rangle: F \times \check{F} \rightarrow R$ is the canonical pairing. The R -module

$$\text{Quat}(F, N) = R \oplus F$$

becomes a quaternion algebra over R under the multiplication

$$(a, u)(b, v) = (ab - \frac{1}{2}N(u, v), av + bu + \frac{1}{2}\varphi_N^{-1}(u \times_\alpha v))$$

for all a, b in R , u, v in F , with norm

$$N_{\text{Quat}(F, N)}((a, u)) = a^2 + N(u).$$

Every quaternion algebra over R is of the type $\text{Quat}(F, N)$. We obtain

$$\text{Quat}(F, N) = (R, F, B, \times_\alpha),$$

where B is the nondegenerate symmetric bilinear form on F associated with N .

We have thus unified the three constructions known for composition algebras.

REFERENCES

- [B] Becker, E., *Über eine Klasse flexibler quadratischer Divisionsalgebren*, J. für Reine u. Angew. Math. 256 (1972), 25-57.
- [E] Elduque, A., *Colour algebras and Cayley-Dickson algebras*, Proc. of the Royal Soc. of Edinburgh, 125A (1995), 1287-1303.
- [K] Knus, M. A., "Quadratic and Hermitian Forms", Springer Verlag, 1991.
- [KOS] Knus, M. A., Ojanguren, M., Sridharan, R., *Quadratic forms and Azumaya algebras*, J. Reine Angew. Math. 303/304 (1978), 231-248.
- [KS] Kunze, R. A., Scheinberg, S., *Alternative algebras having scalar involution*, Pac. J. Math. 124 (1986), 159-172.
- [Ku] Kuz'min, E. N., *A class of anticommutative algebras*, Algebra i Logika 6 (4) (1967), 31-50.
- [M1] McCrimmon, K., *Nonassociative algebras with scalar involution*, Pacific J. of Math. 116(1) (1985), 85-108.
- [M2] McCrimmon, K., "A Taste of Jordan Algebras", Universitext, Springer Verlag, New York 2004.
- [Mo] Morandi, P., *Lie algebras, composition algebras, and the existence of cross products on finite-dimensional vector spaces*, Expo Math. 17 (1999), 063-074.
- [O] Osborn, J. M., *Quadratic division algebras*, Trans. Amer. Math. Soc. 105 (1962), 202-221.
- [P1] Petersson, H.P., *Eine Bemerkung zu quadratischen Divisionsalgebren*, Arch. Math. 22 (1971), 59-61.
- [P2] Petersson, H. P., *Composition algebras over algebraic curves of genus 0*, Trans. Amer. Math. Soc. 337 (1993), 473-491.
- [PR] Petersson, H. P., Racine, M., *Reduced models of Albert algebras*, Math. Z. 223 (3) (1996), 367-385.
- [Pu] Pumplün, S., *Quaternion algebras over elliptic curves*, Comm. Algebra 26 (12) (1998), 4357-4373.
- [T] Thakur, M. L., *Cayley algebra bundles on $\mathbb{A}_K^2$ revisited*, Comm. Algebra 23(13) (1995), 5119-5130.

SCHOOL OF MATHEMATICS, UNIVERSITY OF NOTTINGHAM, UNIVERSITY PARK, NOTTINGHAM NG7 2RD, UNITED KINGDOM, SUSANNE.PUMPLUEN@NOTTINGHAM.AC.UK